ΑD	1		

Award Number: W81XWH-06-1-0191

TITLE: Targeting Androgen Receptor Function by MicroRNA in Prostate Cancer

PRINCIPAL INVESTIGATOR: Girish C. Shukla, Ph.D.

CONTRACTING ORGANIZATION: Cleveland Clinic Foundation,

Cleveland, OH 44109

REPORT DATE: July 2008

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

IT.	EPORT DOC	OMENIAIIO	N PAGE		OMB No. 0704-0188		
Public reporting burden for this	collection of information is esti	mated to average 1 hour per res	ponse, including the time for revie	wing instructions, search	hing existing data sources, gathering and maintaining the		
					llection of information, including suggestions for reducing		
					rson Davis Highway, Suite 1204, Arlington, VA 22202- a collection of information if it does not display a currently		
valid OMB control number. PL	EASE DO NOT RETURN YOU	R FORM TO THE ABOVE ADD	RESS.	g			
1. REPORT DATE		2. REPORT TYPE		3. D	ATES COVERED		
01-07-2008		Annual		1 Ja	an 2007 – 30 Jun 2008		
4. TITLE AND SUBTIT					CONTRACT NUMBER		
4. III EE AND 00D III				Jul.	SONTRAGI NUMBER		
Targeting Androge	n Receptor Function	on by MicroRNA in I	Prostate Cancer	5b. (GRANT NUMBER		
9 9 9	•	,		W8	1XWH-06-1-0191		
					PROGRAM ELEMENT NUMBER		
				36.1	FROGRAM ELEMENT NOMBER		
6. AUTHOR(S)				5d. l	PROJECT NUMBER		
0::1 0 01 11 5	N 5			F	TACK NUMBER		
Girish C. Shukla, F	h.D.			be.	TASK NUMBER		
				5f. V	WORK UNIT NUMBER		
Emaile a shulda@a	anahia adu						
Email: g.shukla@c							
7. PERFORMING ORG	SANIZATION NAME(S)	AND ADDRESS(ES)			ERFORMING ORGANIZATION REPORT		
				N	UMBER		
Cleveland Clinic F	oundation.						
Cleveland, OH 44							
Cievelariu, Ori 44	1						
9. SPONSORING / MC	NITORING AGENCY N	IAME(S) AND ADDRES	S(ES)	10. 9	SPONSOR/MONITOR'S ACRONYM(S)		
U.S. Army Medica	l Research and Ma	teriel Command					
Fort Detrick, Maryl							
Tort Detrick, Mary	and 21702-3012						
				11. \$	SPONSOR/MONITOR'S REPORT		
				l l	NUMBER(S)		
	VAILABILITY STATE						
Approved for Publi	ic Release; Distribu	ıtion Unlimited					
• •	,						
13. SUPPLEMENTAR	VNOTES						
13. 301 I ELMENTAN	INOILS						
	4-4				and a subject of the		
					ancer in North American men and		
the blockade of an	drogen action thro	ugh the AR has bee	en the cornerstone of	f systemic ther	apy of prostate cancer. However,		
the effectiveness of	of this therapy is rat	her transient which	inevitably fails and t	tumor arowth re	esumes despite androgen blockade.		
					s the development of		
androgen-indepen	dent prostate canc	er. Originally we pro	oposed the utilizatior	n of micro (mi)	RNAs to blockade the expression of		
AR in prostate carcinoma cells. We have identified a few miRNAs that can repress the AR protein synthesis in prostate							
carcinoma cells. Our long-term goals are to identify naturally occurring miRNAs that have potential to block the activity of AR							
and to improvise the	neir efficacy by ration	onal designing to pr	ovide novel 'AR Ant	agonist miRNA	NS'.		
15. SUBJECT TERMS							
Anti AR MiRNAs,							
ATTU ATT MITTINAS	Horicoully INIA						
16 SECURITY OF ASS	SIFICATION OF		17 I IMITATION	18 NUMBER	19a NAME OF RESPONSIRI E PERSON		
16. SECURITY CLASS	SIFICATION OF:		17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON		
		I	17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	USAMRMC		
a. REPORT	b. ABSTRACT	c. THIS PAGE					

UU

7

Form Approved

Table of Contents

	<u>Page</u>
Introduction	4
Body	4
Key Research Accomplishments	5
Reportable Outcomes	5
Conclusion	5
References	
Appendices	

INTRODUCTION:

Prostate cancer is the most commonly diagnosed and second most deadly cancer in North American men. Androgen receptor (AR) protein plays an important role in the development and progression of prostate cancer. Blockade of androgen action through the AR has been the cornerstone of systemic therapy of prostate cancer. However, the effectiveness of this therapy is rather transient which inevitably fails and tumor growth resumes despite androgen blockade. The failure of AR receptor antagonists results in higher levels of AR which is one of the causative factors of the development of androgen-independent prostate cancer. We proposed the utilization of micro (mi) RNAs to blockade the expression of AR in prostate carcinoma cells. Our goals are to identify naturally occurring miRNAs that affect the activity of AR and to improvise their efficacy by rational RNA-RNA interactions (between miRNA and 3' mRNA region of AR) to provide novel 'therapeutic miRNAs' which can target the expression of AR.

BODY:


Task 1. To examine if androgen receptor (AR) translation is modulated by a naturally occurring hsa-mir-183 microRNA (miRNA) and to validate that the 3'UTR of AR is a bona fide target of miRNA using chimeric luciferase reporter constructs (Months 1-12).

Task 1a Proposed Work: Design and commercial acquisition and subsequent overexpression of synthetic miRNAs by transient transfections into LNCaP cells and investigation of AR expression levels by Western blot using anti-AR antibodies. Confirmation that hsa-miR-183 is involved in AR translational control will be achieved (Months1-3).

We have cloned genomic regions coding miRNA and putative cis elements into lentiviral vectors as well as in eukaryotic expression vector pcDNA to examine their effect in AR protein repression in prostate carcinoma cells LNCaP (see below).

Task 1b Proposed Work: Development of multiple test and control luciferase reporter expression constructs, each with an engineered 3' untranslated region (miRNA targeted sequences) of AR cloned downstream of the stop codon of luciferase cDNA. Chimeric luciferase and appropriate control reporter constructs will be ready (months 1-6).

We have synthesized the luciferase reporter with androgen receptor 3' UTR cloned in the downstream of the



stop codon. We have chimeric luciferase and it function was tested in transiently transfected cell culture system. Results are shown below:

Figure 1 Testing of firefly luciferase containing 3' UTR of androgen receptor in transiently transfection: Firefly luciferase with or without 3' UTR of AR along with appropriate controls were transiently transfected in CHO cells using polybrene. Cells were DMSO shocked 8 hours posttransfection and cells were harvted at 48 hours. Total protein lysates were prepared and equal amount of proteins from the each transfected wells were used to

perform dual luciferase assay. To repress the expression of firefly luciferase expressing AR 3' UTR we cotransfected pcDNA miR-488. All assays were performed in triplicates and ratio of firefly and renailla luciferase were plotted. pcDNA clone expressing miR-488 was unable to repress the expression of firefly luciferase as shown in "3'UTR+488" in Figure 1. Nonetheless, this expreiment established that our chimeric


firefly luciferase/AR 3'UTR is able to express luciferase in heterologus CHO cell culture systesm. Therefore, we successfully achieved the Task 1b.

Task 1c Proposed Work: Confirmation that AR 3' UTR sequences containing chimeric luciferase reporter constructs are responsive to ectopically expressed synthetic miRNAs. Optimization of reporter assays, evaluation of target specificities and interactions between target sequences and miRNAs will be achieved (Months 6-8).

Currently, we are in the process of optimization of chimeric firefly luciferase and renilla luciferase in order to test the effect of ectopically expressing miRNAs. Also, we are optimizing transfection of pcDNA miR-124a, pcDNA-183 and pcDNA-488 with chimeric firefly luciferase in transient transfection system. In addition, we will generate stably expressing luciferase under Puromycin selection. This would enable us to expedite our transfection experiments.


Additional Work Done: We have generated Hsa-miR-183, Hsa-miR-124a and Hsa-miR-488 expressing lentiviruses for stable expression in LNCaP cells. During this period we tested if stable overexpression of miRNAs in LNCaP cells can target the AR mRNA to modulate the protein synthesis.

We have cloned the coding regions encompassing genomic sequences into Lentiviral vectors. Virus expressing miR-183, miR-124a and miR-488 were generated in 293 T cells. Viruses were purified and concentrated utilizing established protocols. The purified viruses were dissolved in RPMI 1640 medium and LNCaP cells were transduced using these viruses. RNA and protein lysates were produced from these cells to perform qRT-PCR and Western Blot to detect the effect of miRNAs on the RNA and protein levels of Androgen receptor respectively. No significant knockdown of AR protein was observed in LNCaP cells as compared to no virus



lanes. It is possible either these miRNA are not being produced by lentiviruses or AR is not a target of hsa-miR-124 and hsa-miR-183 miRNAs when transfected individually. To check if viral system is producing hsa-miR-124 and hsa-miR-183 expression we performed Northern Blot, however unable to detect the expression of these miRNAs. Ribonuclease protection assays will be done to establish if miRNAs are being expressed from lentiviruses.

Figure 2 A: Western Blot of AR protein prepared from cells expressing miR-124 and miR-83: A time course of 24 hours, 48 hours and 72 hours was done to study the affect of ectopic miR expression on endogenous LNCaP AR. NV- No Virus transduced in LNCaP cells, GFP – cells were transduced with green



fluorescent protein expressing lentivirus, miR-124 and miR-183 corresponds to miRNAs. Beta actin was used as loading control.

Figure 2B: Quantitation of AR protein in lentiviral mediated LNCaP cells: The histogram was plotted on the values of AR protein bands shown in Figure 2a as determined by densitometric scans and image quant program. The miR-124 and miR-183 shows a reduction in AR protein level at 72 hours as compared to 24 hours. Currently, we are in the process of repeating the experiment.




Figure 3a and 3b: These Figures are essentially the similar to experiment shown in Figures 2a and 2b except in this experiment we tested lentiviral miR-488 expression. As shown in the panel 3b '488-72' marked reduction of AR protein was observed. Currently, we are in the process of repeating the experiments.

Figure 3a




Figure 3b

Task 1d Proposed Work: Rational design and testing of designer miRNAs that can modulate the expression of AR 3' UTR containing chimeric luciferase reporter constructs. (Months 8-10).

Work Done: All the experiments stated above will be repeated and a miRNA which can maximally repressed the expression of endogenous LNCaP AR and also represses the firefly luciferase expression in CHO will be rationally 'redesigned'. This will be done by increasing or decreasing the RNA:RNA base-pairing interaction between the 3' AR UTR target and miRNA. So far, miR-488 appears to be a better candidate miRNA that has potential to knockdown the expression of miRNA at greater levels.

Task 2. Proposed Work: Development of an inducible miRNA based cell culture model system to target translational knockdown of endogenous androgen receptor in prostate carcinoma LNCaP cells.

Work Done: Since last report, we got one step closer in the identification of a miRNA which can target the expression of AR. However, we are not able to confirm miRNA mediated effect of AR in cell based assays. Once we identified a miRNA using above mentioned techniques, and identify its role in cellular assays, we will subclone that miRNA in inducible expressing systems in and test for it functionality in the prostate carcinoma cell culture system. This specific aim is depend on the success of Task 1d.

PROBLEMS: The funds from this grant mechanism are ended and it is in "no cost" extension mode. We have applied for more funds from PCRP funding mechanism to continue to work on this project. However, currently we have very limited resource to continue to fund this project. A final report stating all the accomplishment will be submitted in 2009. However, below we describe interim accomplishment of the project.

KEY RESEARCH ACCOMPLISHMENTS: Lentiviral expression vectors expressing miR-124, miR-183 and miR-488 are available. We also have chimeric firefly luciferase with AR 3' UTR construct. As a proof of principle we have tested the expression of chimeric luciferase in CHO cells. The optimization of miR and chimeric firefly luciferase will be pursued, if we are able to generate sufficient funds.

REPORTABLE OUTCOMES: As of now, we do not have reportable outcomes from this funding. However, to continue to build upon the knowledge and resources we have gained during this period, I have applied for New Investigator Award from PCRP.

CONCLUSION: We have developed reporter constructs expressing multiple miRNAs and have opportunity to test for their efficacy in AR inhibition in prostate cancer cells. The experimental identification of miRNAs which maximally repress the translational activity of the AR and further improvisation in their design will allow us to develop a subset of "AR antagonist miRNA".

SO WHAT: The knowledge gained from our efforts so far indicates that our project is heading in right direction. However, we are unable to sustain the momentum due to lack of funds and personnel. We would be able to use our constructs, if resources allow, in blockading the expression of AR in prostate carcinoma cell model systems, and perhaps in mouse model of prostate cancer.